

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
2 June 2005 (02.06.2005)

PCT

(10) International Publication Number  
**WO 2005/049831 A1**

(51) International Patent Classification<sup>7</sup>: **C12N 15/11,**  
15/01

MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC,  
SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA,  
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:  
PCT/SG2003/000274

(84) Designated States (*regional*): ARIPO patent (BW, GH,  
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,  
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(22) International Filing Date:  
20 November 2003 (20.11.2003)

(25) Filing Language: English

(26) Publication Language: English

**Declarations under Rule 4.17:**

— *as to applicant's entitlement to apply for and be granted  
a patent (Rule 4.17(ii)) for the following designations AE,  
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES,  
FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,  
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG,  
MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT,  
RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT,  
TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent  
(BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM,  
ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ,  
TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK,  
EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,  
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG)*

— *of inventorship (Rule 4.17(iv)) for US only*

(71) Applicant (*for all designated States except US*): AGENCY  
FOR SCIENCE TECHNOLOGY AND RESEARCH  
[SG/SG]; 20 Biopolis Way, #07-01, Centros, Singapore  
138668 (SG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): LIU, Ziduo  
[CN/SG]; 30 Medical Drive, singapore 117609 (SG).  
HONG, Yuzhi [CN/CN]; College of Life Science and  
Technology, Huazhong Agricultural University, Wuhan,  
Hubei 430070 (CN). ZHANG, Lianhui [AU/SG]; 30  
Medical Drive, Singapore 117609 (SG).

(74) Agent: LEE, Ai Ming; 80 Raffles Place, #33-00 UOB  
Plaza I, Singapore 048624 (SG).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,

**Published:**

— *with international search report*

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: METHOD

(57) Abstract: A method for enriching the GC base pair content of a DNA molecule the method comprising the steps of (a) providing a DNA template molecule in which at least some of the A residues are base paired with U residues and (b) replicating the DNA template molecule provided in step (a) under conditions in the replication reaction medium in which at least some of the U residues base pair with a G residue. Typically, the DNA template molecule in (a) is produced by replicating a first template DNA molecule in the presence of dUTP so that at least some of the T residues of the first template are replaced by U residues to form a second template molecule. Thus, a preferred method comprises the steps of (1) providing a first template DNA molecule, (2) replicating the first template DNA molecule in the presence of dUTP so that at least some of the T residues of the first template are replaced by U residues to form a second template molecule and (3) replicating the DNA template molecule produced in step (2) under conditions in the replication reaction medium in which at least some of the U residues base pair with a G residue. The invention also includes a method for making a mutant polypeptide with altered properties compared to the polypeptide encoded by a DNA molecule, the method comprising (a) providing a DNA molecule encoding a polypeptide of interest, (b) enriching the GC base pair content of the DNA molecule according to the method set out above, (c) expressing the polypeptide encoded by the DNA molecule whose GC base pair content has been enriched and (d) selecting a polypeptide with altered properties. The invention also includes A mutant A1bD polypeptide wherein Ser40 has been replaced by another amino acid residue.

WO 2005/049831 A1